



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Ontario, a male bird of this species, that showed a strange freak in the tail. While all the other feathers are of normal length and development (2.50 in. long), one shows a length of 3.75 in. It is of the same color as the other tail feathers, but the vanes are narrower. Near the ends of the other rectrices it is much attenuated, as though it had wanted to stop growing at that point. It is the only instance of this kind that has come under my observation.—C. W. G. EIFRIG, *Ottawa, Ont.*

The Migrant Shrike (*Lanius ludovicianus migrans*) at Ottawa, Ont.—This shrike is a common breeder here. In 1904 I saw the first ones, a pair, on April 5, and this year on March 30. They frequent wet meadows, old fields, etc. By the middle of April they are common. The last one in 1904 I saw August 23, when I was attracted to a hedge by the low but pretty singing of a bird, which proved to be the shrike. The song was much like the subdued song of the Catbird, with much of its sweetness. Although those that I took had beetles only in their stomachs, yet on April 26, 1904, I saw a pair that had a Song Sparrow impaled on a thorn and had eaten off the head. May 10 I found the first nest in the usual thorn-thicket in a moist meadow, containing two eggs. May 21, I found another nest at Casselman, thirty miles east of Ottawa; this was eight feet up in a little wild plum tree and contained five young about a day old and one infertile egg. The female, in both cases, would only fly away when the person was within a few feet of the nest, and would sit near by and utter queer rasping or gurgling notes of protest.—C. W. G. EIFRIG, *Ottawa, Ont.*

Capture of the Kirtland Warbler near Richmond, Ind.—On May 13, 1905, a female Kirtland Warbler (*Dendroica kirtlandi*) was shot about six miles east of here by Mr. Loren C. Petry. It was silent, perfectly fearless, and showed the 'teetering' habit after the manner of a Palm Warbler or Titlark. It was in a clump of low bushes.—D. W. DENNIS, *Richmond, Ind.*

The Kentucky Warbler at Winneconne, Wisconsin.—On May 7, 1905, I had the pleasure of finding a Kentucky Warbler (*Oporornis formosa*) in a damp corner of the woods near Lake Winneconne. I observed it for ten or fifteen minutes from a distance of only a few feet, and am thus positive of the identification.

The Kumlien-Hollister List of Wisconsin Birds (1903) says of this species: "Dr. Hoy took one specimen at Racine (May 10, 1851) and we have but six other records for the State for 60 years, all about Lake Koshkonong, in spring."—HENRY P. SEVERSON, *Winneconne, Wisc.*

Wintering of the Brown Thrasher in a Park in New York City.—A Brown Thrasher (*Toxostoma rufum*) passed the winter of 1904-05 in Morningside Park, New York City. I first saw him on Oct. 24, 1904,